



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

compact mass. On the 11th and 12th came freezing weather, and the fall of a small amount of very light snow. On the 13th the thermometer, toward noon, rose above freezing-point, with a stiff breeze from the south. This wind so acted on the surface-particles of the upper layer of uncompacted snow as to set some of them in motion. Each particle thus set in motion, owing to the soft condition of the surface-snow, formed a nucleus, which, as it proceeded, forced along by the wind, gathered the contiguous portion of the soft layer, and assumed the form of a cylinder, with a conical cavity at each end, and having a length about twice as great as its diameter. The size depended upon the inclination and smoothness of the surface traversed. The largest cylinders I saw were about three feet long, at which limit they acquired sufficient weight to indent the frozen surface of the under or main body of the snow. This, of course, stopped the further rolling of the mass. The only locality where they acquired the above large size was where the surface had a slight inclination to the north; and the snow was deep enough to cover all weeds, leaving a perfectly uniform and smooth surface for their formation. In many cases the balls were rolled up an inclination of as much as one foot in ten, when exposed to the unbroken force of the wind; but those thus formed acquired weight sufficient to resist the pressure of the wind, when about six inches in diameter. When the surface inclined toward or directly away from the wind, the balls traversed a straight path; but, when the surface declined to the north-east or north-west, the path was a curve; at its initial, approximately straight; but, as the ball acquired weight, its direction was a compromise between that required by gravity and that by the direction of the wind, until, in some cases, the ball obeyed gravity alone. The most curious part of the display was the abundance of the balls. While travelling three miles, I saw what I estimated at over a hundred acres dotted more or less thickly with the cylinders. In some cases there were twenty-five balls to the square rod; in others, only two or three; averaging, perhaps, eight or ten. I saw multitudes in the process of formation, which was as sudden as a flash; but they almost immediately assumed a slow rate of motion, about that of a mole taking his leisurely walk. In a few cases the cylinders would stop, and afterward be forced into motion again. The largest examples required for their formation the traversing of from two to three rods. SAM HUSTON.

Richmond, O., Jan. 16, 1884.

The wind performed a very pretty feat in some portions of northern Ohio on the morning of Jan. 13. Loose bits of snow were caught up as a nucleus, and rolled along upon the surface until balls of considerable size and peculiar shape were formed. The whole surface was strewn with the balls; but they were most abundant upon lawns and fields where the wind was not obstructed, every square yard, in some places, bearing a ball of greater or less size. The largest observed here were upon the college ball-grounds, where they reached ten inches in height, and a horizontal length of eighteen inches. Even these were swaying as the gusts passed over them; and their tapering track could be plainly traced back towards the south-west, twenty-five or thirty feet, to the apex where they started. Their shape was cylindrical, deeply hollowed at both ends, so that they looked like 'muffs,' and the spiral formed by the successive layers was finely regular and distinct.

The meteorological conditions which made the phe-

nomenon possible were as follows. Two days before the occurrence a slight crust was formed upon the snow. On the following day an inch of light flaky snow fell upon this crust. Then followed the warm south-west wind on the morning of the 13th, which brought the upper layer of snow into the adhesive state, and rolled the balls before the crust was weakened; the crust sustaining the balls, and keeping them up to the wind, and at the same time furnishing a smooth floor upon which they could be propelled. The nuclei of the balls were obtained from chance foot-tracks, walk-borders, lumps blown from trees, etc., though often it was difficult to account for them. The balls were most abundant and perfect at about nine o'clock A.M. Before noon the crust had been attacked, and all sunk to rounded, insignificant clumps. Oberlin, O. ALBERT A. WRIGHT.

[Similar snow-rolls were seen at Sharpsville, Mercer county, Penn., on the same day, by J. M. Goodwin.]

Halos round the moon.

On the evening of Jan. 12, at 9.20 (90th meridian time), my attention was called to a peculiar appearance about the moon. The sky was quite clear at the time, and there appeared around the moon several colored circular bands. The first was of a bright silver-gray shade, and about two diameters of the moon in width. The next was yellow, the next faint orange, and the next violet. The three bands were each about one-half a diameter in width. The outermost band was of a green shade, and about two diameters in width. At ten o'clock the innermost light band remained, but all the others had been replaced by a blue band lighter than the surrounding sky.

H. A. HUSTON.

Lafayette, Ind., Jan. 14, 1884.

Explorations in Guatemala.

Looking over the back numbers of your esteemed journal, I came across a slight error. In the article 'Lorillard City' it is said (ii. 412), "M. Charnay found the ruins of an ancient city, which he named after his generous patron. In his exploration here, he was assisted by a young Englishman, Mr. Alfred Maudslay, with whom he shares the honor of discovery," etc.

Neither Mr. Maudslay, who arrived at these ruins before Mr. Charnay, nor the latter, can claim this honor. In fact, Mr. Maudslay distinctly states (p. 196 of the *Proc. roy. geogr. soc.*, April, 1883) that they have been discovered by Mr. Edwin Rockstroh, tutor on the Lyceo nacional at Guatemala City. This gentleman made, during the first half of 1881, a geographical and archeological exploration in the northern and western parts of the republic, visiting Tikal, and navigating the Río de la Pasión, Río de las Salinas, Río de los Gacandones, and the Usumasinta as far down as the ruins mentioned. He sent a short account of this voyage to Petermann's *Mittheilungen* (1881, p. 396).

In that account Mr. Rockstroh mentioned particularly the building described by Mr. Maudslay on p. 198 of the geographical society's proceedings; and (1882, on p. 435) he clearly states that Charnay's 'Lorillard City' is the same as that discovered by him in 1881. Mr. Rockstroh mentioned in his first letter to the *Mittheilungen* (July 19, 1881), that the Gacandones call these ruins 'Menche,' and promised in his last notice (1882, p. 435) an explanation of this name. I am not aware that he has furnished one.

I find in the 'Historia de la provincia de San Vicente de Chiapa y Guatemala,' by Antonio de